

REPORT DOCUMENTATION PAGE			Form Approved OMB NO. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comment regarding this burden estimates or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.				
1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE January 26, 1999		3. REPORT TYPE AND DATES COVERED Final 6/1/95-11/30/98
4. TITLE AND SUBTITLE Designed Synthesis and Assembly of Molecular Building-Blocks for New Materials			5. FUNDING NUMBERS DAAHO4-95-1-0145	
6. AUTHOR(S) Russell N. Grimes				
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) University of Virginia Charlottesville, Virginia 22901			8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Research Office P.O. Box 12211 Research Triangle Park, NC 27709-2211			10. SPONSORING / MONITORING AGENCY REPORT NUMBER ARO 34065.15-CH	
11. SUPPLEMENTARY NOTES The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision, unless so designated by other documentation.				
12a. DISTRIBUTION / AVAILABILITY STATEMENT Approved for public release; distribution unlimited.			12 b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words)			19990616 161	
<p>The project centered on the designed synthesis of small metallocarborane sandwich complexes for use as building-blocks in the construction of new families of electroactive, magnetoactive, and/or optically active materials. The compounds of interest exhibit generally high thermal and oxidative stability, chemical versatility, electronic tunability, and solubility in organic solvents. Synthetic routes were developed for several new classes of multi-metallic metallocarborane clusters that are bifunctional and suitable as precursors to novel metallopolymers. These include B-B, B-X-B (X = alkene, alkyne, or aryl), and M-M linked C₂B₄M clusters, arene- and carborane-endcapped multidecker sandwiches, and B-halo species. Strategies for preparing high molecular weight electronically active materials from these compounds were explored. Concurrently with the synthetic work, the electronic structures, nonlinear optical properties, and other aspects of selected systems were investigated in detail, partly in collaboration with other laboratories.</p>				
14. SUBJECT TERMS carboranes metallocarboranes cobaltacarboranes			15. NUMBER OF PAGES 9 + ii	
metal-arene complexes sandwich complexes multidecker sandwich complexes			16. PRICE CODE	
17. SECURITY CLASSIFICATION OR REPORT UNCLASSIFIED			18. SECURITY CLASSIFICATION OF THIS PAGE UNCLASSIFIED	
19. SECURITY CLASSIFICATION OF ABSTRACT UNCLASSIFIED			20. LIMITATION OF ABSTRACT UL	


I. Statement of Problem

The main objective of this work was to design and synthesize small metallocarborane sandwich complexes that are suitable as building-blocks in the construction of new families of electroactive, magnetoactive, and/or optically active materials, and to conduct detailed studies of the electronic and molecular structure of selected systems of this class.

II. Summary of Important Results


1. Synthesis of Heavy Transition-Metal Metallocarborane Sandwich Complexes

Synthetic routes to multidecker sandwich complexes of second- and third-row transition elements were developed. As an example, double-decker iridium and ruthenium complexes were prepared and employed as synthons for preparing several new classes of triple- and tetradeckers, including rhodium- and iridium-centered sandwiches including compound 9 ("Group9ocene"), a unique sandwich containing all three Group 9 metals. (For details see Interim Technical Report for February 1996, and publication 9 cited below).





2. Cluster Linkage.

In a project begun during the previous 3-year grant and extended under the current project, we developed methods for linking metallocarborane clusters via B-B, C-C, and metal-metal (M-M) intercluster bonds, for the purpose of eventually building extended electron-delocalized solid-state metallocarborane-based networks. Paper 4 cited below describes the B-B and C-C linking reactions, both of which are novel to boron chemistry:




3. Cage Recapitulation.

A further dimension to cluster linkage methodology was added via "recapitation", a process that is functionally the reverse of our "decapitation" reaction in which the apex BH in a closo- MC_2B_4 cage is removed to generate an open face which in turn can π -complex to metal ions (e.g., conversion of **3a-d** to **4a-d** shown above). In recapitation, illustrated here, a boron atom bearing a substituent is inserted into the open face to a B-X unit, thereby affording apex-substituted clusters that can be used to form apex-linked oligomers and polymers (paper 10). In collaborative work with Siebert and co-workers at the University of Heidelberg (paper 14), this approach was exploited to prepare and structurally characterize a series of linked biscobaltacarboranes:




4. Metal Hydride Sandwich Complexes.

Manipulation of metal-bound protons in hydridometallacarboranes of iron and cobalt (e.g., their reversible deprotonation with accompanying redox processes at the metal centers) was explored in our laboratory as a means of controlling metal oxidation states. This approach has potential as a type of "molecular switch" that could be incorporated in organometallic polymers to alternate between conducting and insulating behavior. As described in paper 12, this work has been extended to ruthenium hydrides with the synthesis of the double-decker species $\text{Cp}^*\text{Ru}^{\text{II}}\text{H}(\text{Et}_2\text{C}_2\text{B}_4\text{H}_4)$, which undergoes an unexpected B-alkylation on treatment with methyl triflate:




A heterobimetallic RuH-Co triple-decker was also synthesized as shown.




5. Metal Carbonyl-Based Chemistry and Metal-Metal-Bonded Clusters.

A different approach to metallocarborane sandwich synthesis, with implications for materials engineering, utilizes metal carbonyl complexes whose CO ligands are more easily displaced than hydrocarbon ligands such as Cp, Cp*, and arenes, and hence promote coupling reactions directly at the metal centers. The following schemes summarize our work on molybdenum and tungsten metallocarboranes which has generated some novel species including the first M-M linked small metallocarboranes (paper 15). X-ray crystallography has established several of the structures shown, including the bent tetradecker (the first of its genre in carborane chemistry).




The principal significance of this work is twofold: (1) double-open-ended complexes such as the Mo_2 species shown, are potential precursors to linear M-M linked polymers which would be quite novel and may have interesting electronic and magnetic properties; (2) the absence of hydrocarbon ligands makes these species attractive candidates for conversion to graphite-free metal-boron solid-state materials.




6. Double-Decapped Metallacarborane Synthons Complexes.

After an extended effort, we have found a way to prepare small double-decker complexes containing *two* open-faced C_2B_3 rings that can be deprotonated and reacted with metal ions to generate extended multidecker sandwiches, i.e., polydeckers. Such species have been a goal of this research for some time, and we are currently investigating the actual formation of such polydecker species.



7. Alkyne-Linked Systems.

Substantial progress has been achieved in the synthesis of metallacarboranes containing multiple metal centers with alkyne (bare carbon atom) connectors that are efficient conductors of electrons and are often called "molecular wires". The double-open-ended complexes shown are well positioned to serve as synthons for alkyne-linked conducting polymers, a prospect we are currently investigating.



III. Publications produced under this Contract

- (1) "Cluster Forming and Cage Fusion in Metallocarborane Chemistry", Grimes, R. N., *Coord. Chem. Rev.* **1995**, *143*, 71.
- (2) "Organotransition-Metal Metallocarboranes. 39. Arene-Capped Ruthenium-Carborane Tetradecker Sandwich Complexes", Greiwe, P.; Sabat, M.; Grimes, R. N., *Organometallics* **1995**, *14*, 3683.
- (3) "Organotransition-Metal Metallocarboranes. 40. Regiospecific Halogenation of Transition Metal Small Carborane Complexes", Stockman, K. E.; Garrett, D. L.; Grimes, R. N., *Organometallics* **1995**, *14*, 4661.
- (4) "Organotransition-Metal Metallocarboranes. 41. Synthesis and Structure of B-B and Cp*-Cp* Linked Cobaltacarborane Clusters", Wang, X.; Sabat, M.; Grimes, R. N., *Organometallics* **1995**, *14*, 4668.
- (5) "Organotransition-Metal Metallocarboranes. 42. Synthesis and Cluster Fusion of Iron-Centered Tetradecker Sandwiches", Wang, X.; Grimes, R. N., *Inorg. Chem.* **1995**, *34*, 6509.
- (6) "Organotransition-Metal Metallocarboranes. 43. Directed Synthesis of Carborane-Endcapped Multidecker Sandwiches", Wang, X.; Sabat, M.; Grimes, R. N., *J. Am. Chem. Soc.* **1995**, *117*, 12218.
- (7) "Organotransition-Metal Metallocarboranes. 44. Synthesis and Structure of Pentadecker and Hexadecker Sandwiches", Wang, X.; Sabat, M.; Grimes, R. N., *J. Am. Chem. Soc.* **1995**, *117*, 12227.
- (8) "Metal-Carborane Multidecker Sandwich Complexes as Building-Blocks for New Materials", Grimes, R. N., *Applied Organometallic Chemistry.*, **1995**, *10*, 209-225.
- (9) "Organotransition-Metal Metallocarboranes. 46. Multidecker Sandwiches of the Group 9 Metals", Franz, D. A.; Houser, E. J.; Sabat, M.; Grimes, R. N., *Inorg. Chem.* **1996**, *35*, 7027.
- (10) "'Recapitulation' of Nido-Metallocarboranes as a Synthetic Tool: Preparation of Apically Substituted CoC₂B₄ Clusters via Boron Insertion", Curtis, M. A.; Sabat, M.; Grimes, R. N., *Inorg. Chem.* **1996**, *35*, 6703.
- (11) "Metal-Carborane Sandwiches and Macromolecular Assemblies", Grimes, R. N., in *Advances in Boron Chemistry*; Siebert, W., Ed.; Royal Society of Chemistry: Cambridge, U.K., 1997, p. 321-332.
- (12) "Organotransition-Metal Metallocarboranes. 45. Synthesis and Structure of Hydridoruthenacarborane Double-Decker and Triple-Decker Sandwich Complexes", Houser, E. J.; Curtis, M. A.; Sabat, M.; Grimes, R. N., *J. Organometal. Chem.* **1996**, *536/537*, 115.

- (13) "Recent Advances in Metallocarborane Sandwich Chemistry: Controlled Linking, Stacking, and Cool Fusion", Grimes, R. N., in *The Borane-Carborane-Carbocation Continuum*, Casanova, J., Ed.; Wiley-Interscience, **1998**, Chapter 13, pp. 307-329.
- (14) "Apically Linked Small Metallocarborane Clusters. Directed Synthesis and Structural Characterization of $7,7'-[\text{Cp}^*\text{Co}(2,3\text{-Et}_2\text{C}_2\text{B}_4\text{H}_3)_2]$ and $[\text{Cp}^*\text{Co}(2,3\text{-Et}_2\text{C}_2\text{B}_4\text{H}_3-7)]_2\text{X}$ Complexes ($\text{X} = \text{MeCH}$, $\text{HC}=\text{CH}$, or $\text{C}\equiv\text{C}$)", Curtis, M. A.; Müller, T.; Beez, V.; Pritzkow, H.; Siebert, W.; Grimes, R. N., *Inorg. Chem.* **1997**, *36*, 3602.
- (15) "New Metallocarborane Chemistry: Molybdenum and Tungsten Carbonyl Multidecker Sandwiches. Cluster Dimers Linked via Metal-Metal Bonds", Curtis, M. A.; Houser, E. J.; Sabat, M.; Grimes, R. N., *Inorg. Chem.* **1998**, *37*, 102.
- (16) "A Delocalized Analogue of the Bicobaltocene Cation Derived From Reduction of a d^6d^6 (Fulvalenediyl)bis(cobaltacarborane) Complex: Electrochemistry, ESR, Optical Spectroscopy, and IR Spectroscopy of the Carborane Group", Chin, T. T.; Grimes, R. N.; Geiger, W. E., *Inorg. Chem.* **1999**, *38*, 93.

Manuscript in press:

- (17) "Small Carborane Ligands as Spectators and as Players", Grimes, R. N., *J. Organometal. Chem.*, in press.

Manuscript submitted:

- (18) "A Novel Heterooctametallic Metallocarborane Tetramer", Parker, K. E.; Russell, J. M.; Sabat, M.; Grimes, R. N., submitted.

IV. Participating Scientific Personnel

"Studies in Metallocarborane Chemistry: Controlled Cluster Synthesis and Reagents for Organic Transformations", M. A. Curtis, Ph. D. University of Virginia, 1998

Principal Investigator: Russell N. Grimes, Professor of Chemistry

Postdoctoral Associates:

Keith Parker
Dennis Malaba

Graduate Students (degrees earned):

Michael Curtis (Ph.D. 1998)
Barbara Gilmore
Dana Kobeda

V. Inventions: none